


Journal Pre-proof

Facilitating Thought Progression: A Neurocognitive Framework Linking Thought Dynamics and Mood Disorders

Moshe Bar, Shira Baror



PII: S0006-3223(26)01219-9

DOI: <https://doi.org/10.1016/j.biopsych.2026.04.018>

Reference: BPS 16105

To appear in: *Biological Psychiatry*

Received Date: 7 January 2026

Revised Date: 6 April 2026

Accepted Date: 24 April 2026

Please cite this article as: Bar M. & Baror S., Facilitating Thought Progression: A Neurocognitive Framework Linking Thought Dynamics and Mood Disorders, *Biological Psychiatry* (2026), doi: <https://doi.org/10.1016/j.biopsych.2026.04.018>.

This is a PDF of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability. This version will undergo additional copyediting, typesetting and review before it is published in its final form. As such, this version is no longer the Accepted Manuscript, but it is not yet the definitive Version of Record; we are providing this early version to give early visibility of the article. Please note that Elsevier's sharing policy for the Published Journal Article applies to this version, see: <https://www.elsevier.com/about/policies-and-standards/sharing#4-published-journal-article>. Please also note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2026 Published by Elsevier Inc on behalf of Society of Biological Psychiatry.

Title Page

Title: Facilitating Thought Progression: A Neurocognitive Framework Linking Thought Dynamics and Mood Disorders

Authors:

Moshe Bar^{1,2}

Shira Baror^{3,4}

¹ The Gonda Multidisciplinary Brain Research Center, Bar-Ilan University, Ramat Gan 52900, Israel

² Department of Psychiatry, Massachusetts General Hospital, Boston MA 02114

³The Edmond and Lily Safra Center for Brain Sciences, Hebrew University of Jerusalem, Israel

⁴ Department of Psychology, Max Planck Institute, Leipzig, Germany

Corresponding Author:

Moshe Bar, mbar@mgh.harvard.edu

Running title:

Facilitating Thought Progression

Keywords:

Mood disorders, State of mind, Creativity, Depression, Thought disorders, Inhibition

Abstract

Mood and thought are tightly coupled, but the mechanisms linking them are not understood. This link is particularly important when considering mood disorders such as depression. We propose the Facilitating Thought Progression (FTP) framework, which characterizes depression as a disorder of thought dynamics, encompassing both the temporal evolution and semantic expanse of mental activity. Five parameters jointly determine the fluency of thought progression: breadth, speed, flexibility, novelty, and scope. Diminished progression, reflected by repetitive, slow, and narrowed thinking, is associated with excessive inhibition, reduced plasticity, blunted reward signalling, and overly deep attractor dynamics. Conversely, interventions that enhance associative expansion, cognitive flexibility, and exploratory behaviour restore adaptive brain dynamics. FTP reframes depression as a disorder of cognitive flow rather than content, and provides a mechanistic, testable model for translational research.

Main Text

Introduction

Major depressive disorder (MDD) and related mood disorders are among the most disabling conditions worldwide, affecting over 280 million individuals and contributing substantially to global disease burden (1). Mood shapes nearly every domain of psychological functioning: it influences perception (2), memory (3, 4), attention (5), motivation (6), decision-making (7), physical health (8, 9), and vulnerability to mental disorders (10). Yet despite advances in pharmacological and psychotherapeutic treatments, the neurobiological mechanisms that give rise to core depressive symptoms such as psychomotor slowing, cognitive rigidity, or anhedonia remain incompletely understood.

A central but underexamined feature of depression is that it alters how thoughts unfold over time. Thinking is inherently dynamic: the mind progresses from one concept to another through a semantic network of memory representations (11). Thinking can advance quickly, traverse broad associations, and flexibly shift perspectives. Alternatively, thinking can be slow or remain confined to a narrow thematic region (12-14). Depression is consistently marked by ruminative, repetitive, and sluggish thinking (15). While extensive research has focused on the *content* of those negative thoughts (14, 16-19), far less attention has been allocated to the possibility that the *dynamics* of thinking play a role in mood.

Growing evidence supports this view. Neural studies show that depression is associated with reduced variability and altered transitions in large-scale networks, including the default mode, salience, and frontoparietal networks (15, 20). Recent dynamic modelling of spontaneous thought further suggests separable clusters including 'stuck' patterns versus more flexible mind-wandering, consistent with an emphasis on trajectories rather than content alone (21).

Here, we propose the *Facilitating Thought Progression (FTP) framework*, a mechanistic model linking mental flow and mood through convergent neurocognitive paths. FTP identifies five dimensions that jointly determine how fluidly the mind traverses the representational landscape. Each dimension reflects a neurocognitive process that may be altered in depression. Below, we describe each dimension and discuss how their modulation may restore adaptive mood dynamics.

Dimensions of thought progression and their relation to mood

Breadth: Associative Expansion and Network Connectivity

Breadth refers to the semantic distance between successive thoughts. Positive mood reliably increases associative breadth, enabling remote associations and recruitment of broader semantic fields (22-25). Behaviourally, broadened states display more unusual word associations. In contrast, negative mood constrains associative spread, producing short-range, repetitive transitions that characterize ruminative thought (26, 27). Depressed individuals often cycle through a narrow cluster of representations with minimal semantic distance, a hallmark of maladaptive spontaneous thought (28). Longitudinal evidence indicates that this narrowing of associative trajectories predicts later depressive symptoms (29-31), suggesting a mechanistic role in mood deterioration. The link between breadth of thought and mood is bidirectional, as experimental manipulations that increase associative breadth causally improve mood (32-34). Semantic priming provides a complementary assay of associative dynamics by indexing spreading activation. In depression, priming effects are often altered, but vary with task parameters (e.g., automatic vs. controlled retrieval), consistent with state-dependent changes in the efficiency of associative propagation rather than a content-specific effect (35). Neuroimaging work further indicates that depression alters the neural implementation of priming (36). Importantly, priming can shift with clinical improvement, suggesting partial normalization of associative dynamics with recovery (37). Although relatively few studies have tracked semantic priming across antidepressant treatment, antidepressants reliably modify emotional information processing, suggesting that pharmacotherapy could alter priming-related indices of associative propagation (38, 39).

At the neural level, reduced breadth corresponds to lower variability and diminished functional integration within the default mode network (DMN) and weakened coupling with frontoparietal systems (20). Such reduced network flexibility limits the brain's ability to traverse distant semantic spaces, reinforcing the behavioral narrowing seen in depression.

Clinical interventions, including psychedelic-assisted therapy, have been shown to enhance large-scale network integration and reduce depressive symptoms, in line with broadening associative trajectories (40).

Speed: Temporal Dynamics of Neural Processing

Speed reflects the temporal rate at which thoughts evolve. Behaviourally, depressed individuals exhibit decreased thought speed (41-44), and broader deficits in processing speed (45, 46). These cognitive signatures parallel psychomotor retardation (47), fatigue (48) and diminished motivation (49), highlighting speed as a cross-cutting dimension of depressive impairment.

Depression is associated with slowed cortical dynamics and reduced metastability, an index of how rapidly the brain shifts between transient functional configurations (15). Mechanistically, slowed thinking has been linked to an imbalance in cortical excitation-inhibition dynamics and alterations in thalamocortical oscillatory activity, both of which constrain neural transition rates (15, 50).

Clinically, accelerating cognitive tempo through rapid reading or thought-acceleration paradigms reliably elevates mood (51-53). Interventions that enhance fast-frequency neural activity, such as gamma-range neuromodulation, likewise increase mood and cognitive fluency by restoring the rate of neural transitions.

Flexibility: Cognitive Switching and Neural Reconfiguration

Flexibility refers to the ability to shift between mental sets, perspectives, or strategies, and depends on the capacity of large-scale neural networks to reconfigure efficiently. Depression is marked by reduced cognitive flexibility, evident in impaired task-switching (54), lower divergent and convergent creative output (23, 32, 55-57), and difficulty disengaging from habitual thought patterns (58, 59). Importantly, cognitive inflexibility prospectively predicts the onset of depressive episodes, highlighting flexibility as a vulnerability factor (59, 60).

At the neural level, flexibility is supported by functional diversity within the frontoparietal control network (61, 62) and by reconfiguration of prefrontal-hippocampal circuits (63, 64). Depression is associated with reduced network diversity and diminished synaptic adaptability, limiting the system's ability to shift between cognitive modes (65, 66). These constraints map directly onto behavioral inflexibility: when networks cannot disengage from a current state, thought becomes repetitive and rigid.

Pharmacological interventions that enhance plasticity provide evidence for a causal link between neural reconfiguration and cognitive flexibility. Psilocybin increases both neural and cognitive flexibility in depressed patients through serotonergic and network-level mechanisms (67), while ketamine promotes rapid synaptogenesis and dendritic growth, enabling new cognitive trajectories (65). These findings converge with behavioral work showing that inducing creative thinking (32, 68) or increasing task-switching demands improves mood by facilitating exits from maladaptive cognitive loops.

Novelty: Reward-Driven Exploration and Cognitive Updating

Novelty reflects mentally visiting less-frequented cognitive paths and it engages neural systems that promote exploration. Depressed patients exhibit compromised processing of novel stimuli and a tendency to return to familiar cognitive themes (69-71).

Novel information elicits phasic dopaminergic responses in the mesolimbic system, particularly the ventral tegmental area (VTA) and nucleus accumbens, linking exploratory cognition to intrinsic reward (72, 73). These signals facilitate shifting away from overlearned representations and support the updating of internal models, making novelty a key driver of cognitive progression. In depression, this novelty-reward linkage is impaired. Blunted dopaminergic signalling and reduced reward sensitivity diminish the motivational value of novel information (74, 75). Consequently, thought patterns fail to activate reward circuitry, depriving cognition of the intrinsic reinforcement that normally encourages exploratory transitions. Consistent with this

view, real-world indices of exploration (e.g., variability in daily movement patterns) are found to relate to affect and social connectivity, supporting links between exploratory behaviour and well-being (76).

Interventions that re-engage novelty processing may restore exploratory thought. However, chronic engagement in ruminative, repetitive thought has been associated with structural alterations in cortical volume (77), suggesting that reversing entrenched cognitive patterns may require sustained effort and extended training.

Scope: Local-Global Balance in Cognitive Integration

Scope refers to whether attention and thought are locally or globally focused. Depression is characterized by a local attentional scope (78) and the *attentional scope model of rumination* proposes that such narrowing reduces the likelihood that alternative or positive representations enter awareness (79). In contrast, positive mood broadens attentional and conceptual scope, increasing the probability that new information will be incorporated. This broadening effect has been demonstrated across multiple domains, including attentional selection (80, 81), perceptual processing (16), associative breadth (82), abstract construal (83), and behavioral action tendencies (84).

At the neural level, narrowed scope reflects reduced top-down modulation from frontoparietal and fronto-cingulate control regions, which impairs global integration and promotes a local, self-focused processing style (85-87). This diminished control allows DMN dominance and restricts transitions to attentionally broader, contextually informed representations.

Clinically, these mechanisms align with therapeutic strategies that explicitly aim to expand perspective and broaden cognitive scope, including metacognitive therapies, reappraisal-based approaches, and attentional broadening interventions (88, 89). Biological interventions reinforce this principle: psilocybin treatment has been associated with enhanced large-scale network integration and increased coherence across distributed cortical systems (40), changes that likely facilitate global information integration.

Interaction and Dissociability of FTP dimensions

The five dimensions defining the multiscale landscape of thought progression are distinct yet somewhat interdependent: for example, reduced speed can constrain associative breadth, and impaired flexibility might amplify the effects of excessive inhibition on scope. At the same time, interdependence does not imply redundancy. In an integrated cognitive system, the dimensions remain conceptually and computationally distinguishable and can, in principle, vary independently, even if they often covary clinically. Because they interact, empirical progress requires definitions that permit dissociation. Breadth indexes how far successive thoughts travel in semantic space; Speed indexes the rate of transitions; Flexibility indexes the probability of

switching away from an active attractor, goal, or cognitive set; Novelty indexes the selection of low-frequency, high-information-gain states; and Scope indexes the integration window or level of abstraction, which can vary even when semantic distance does not (Fig. 1). Furthermore, FTP does not assume that more is always better: each dimension likely relates non-monotonically to adaptive functioning, with an optimal balance between exploration and stability. Candidate behavioral assays, computational metrics, neural readouts, and measurement guidance are summarized in Supplementary S1.

Figure 1

Figure 1. Relating the dimensions of FTP and mood: A schematic radar-plot illustration of the proposed dimensions of thought progression, with potential individual profiles of positive mood (light blue), neutral mood (blue) and negative mood (dark blue). Different combinations of these thought dimensions are expected to yield different affective states.

Mechanisms Linking Thought Dynamics and Mood

Cortical Inhibition and the Constriction of Thought Flow

Depression is associated with dysregulated prefrontal control, often manifesting as inefficient or reduced recruitment of dorsolateral prefrontal cortex (dlPFC) during demanding cognitive control tasks (90, 91). In other contexts, particularly those involving negative self-referential processing, compensatory control, or rumination, depression can instead be associated with increased prefrontal engagement (92). These context-dependent alterations can yield excessive top-down inhibition on associative and mnemonic systems (including hippocampus and medial temporal cortex (50, 93, 94)), thereby narrowing the progression of thought and increasing the tendency to remain within familiar self-referential basins (94).

In computational terms, heightened inhibition reduces the transition probability between thoughts, leading to cognitive inertia and perseverative loops (95). Conversely, interventions that regulate cortical inhibition, such as serotonergic psychedelics, transcranial magnetic stimulation

(TMS), or GABAergic modulation, can restore thought progression by increasing neural entropy and enhancing connectivity between large-scale networks (40, 96). Psilocybin, for example, has been recently found to reduce rumination and thought suppression in depressed patients (97). Along these lines, evidence from creativity research suggests that reduced inhibition in combination with broader associative activation enables more flexible and creative thought (70, 98, 99). This highlights the role of inhibitory mechanisms in regulating the breadth and fluidity of thought, such that excessive inhibition constrains associative movement while moderate inhibition supports exploratory cognitive trajectories.

Note that within the FTP model, a moderate level of inhibition should be beneficial for stabilizing thought while allowing flexible transitions. Both extremes, hyper-inhibition (ruminative stagnation) and hypo-inhibition (disorganized or psychotic thought), represent departures from optimal thought progression (100, 101).

Cognitive load and Resource Bottlenecks

While excessive inhibition constrains thought through increased top-down suppression, cognitive load restricts thought progression through resource depletion. These mechanisms can co-occur but represent distinct pathways to cognitive stagnation. Rumination imposes an endogenous cognitive load that draws heavily on working-memory and executive-control systems, limiting the capacity for representational updating (102, 103). Neuroimaging work demonstrates that depression is associated with dlPFC–parietal resource depletion and reduced prefrontal efficiency during demanding tasks (2, 104), consistent with impaired gating of working-memory representations as described in computational models of control. Along similar lines, extended durations of depleted resources induce mental passivity (105). High cognitive load narrows associative breadth and reduces the spread of activation (106, 107), thereby constraining thought progression even when inhibition is not elevated. In this context, negative thoughts repeat not because of their content, but because the system lacks the resources to progress beyond them.

Sleep disturbance is likely to exacerbate this resource-bottleneck pathway. In depression, insomnia and fragmented sleep are associated with poorer cognitive performance, including executive function and processing speed (108, 109). Experimental acute sleep loss and fatigue studies further indicate similar reductions in control efficiency and increased perceived effort (110, 111), thereby lowering the capacity to sustain adaptive transitions in thought. Within FTP, sleep-related depletion is expected to manifest as longer dwell times, reduced transition rates, and narrower semantic trajectories.

Attractor Dynamics and Reduced State Transitions

A complementary computational account of thought stagnation comes from attractor-network models of cognition. When inhibition is high or network flexibility is reduced, thought tends to

settle into habitual attractor states of repetitive patterns (112). This is consistent with the constructed theory of emotion, which proposes that the depressed mind fails to effectively update internal models in response to new information (113), as well as recent computational accounts suggesting that mood disorders are marked by low transition probabilities between top-down and bottom-up processing states (95). Empirically, negative mood has been shown to hinder learning from new experiences (114), and the States of Mind framework similarly posits that healthy cognition depends on fluid transitions between internal states that are disrupted in depression (115). Crucially, attractor dynamics represent an emergent systems-level property of recurrent network interactions, and therefore cannot be reduced to excessive inhibition or cognitive load; rather, they reflect a distinct mechanism through which depression deepens basin stability, increases dwell time in maladaptive states, and restricts transitions through thought space. Figure 2 schematizes this contrast by illustrating a more traversable landscape in which thought moves more broadly across states versus a deeper, dominant attractor in which thought becomes stuck for longer periods.

Figure 2

Figure 2. Schematic attractor landscapes and state visitation in thought progress versus thought stagnation: (A) Conceptual thought landscapes in a two-dimensional state space. The left panel shows a more traversable landscape with shallower attractors, whereas the right panel shows a depression-like landscape with a deeper dominant attractor, reflecting greater trapping of thought. (B) Corresponding trajectory-based state visitation. Shading indicates log visit probability, and H indicates normalized Shannon entropy, with lower values reflecting more concentrated time in fewer states and higher values reflecting broader exploration. *Occupancy maps were derived from simulated noisy gradient-based trajectories, followed by spatial binning, smoothing, normalization, and log transformation. Axes represent abstract thought-state dimensions.

Reward Deprivation and the Cost of Stagnation

Depression is marked by profound disruptions in reward processing, reflected in reduced dopaminergic transmission within the mesolimbic system, including the ventral tegmental area, nucleus accumbens, and orbitofrontal cortex (116). FTP proposes that mental progression itself is intrinsically rewarding: traversing novel or richly connected associative paths engages dopaminergic and opioidergic systems, generating hedonic value (117-119). This view aligns with evidence that information seeking, curiosity, and epistemic surprise are rewarding, even without external reinforcement (120-122). Importantly, the hedonic value of richly connected information declines under cognitive load (123, 124) and repeated information is frequently experienced as aversive (125, 126), suggesting that novelty and associative richness are central drivers of intrinsic reward.

Opioidergic mechanisms may also contribute. Mu-opioid receptor density increases along the visual cortical hierarchy, implying that deeper and more elaborated perceptual processing triggers greater endorphin release (127). Dysregulated opioid activity in depression (128) may therefore diminish the hedonic impact of cognitive exploration and contribute to reduced novelty seeking (107, 129).

Critically, FTP frames reward deprivation, reduced exploration, and thought rigidity as a mutually reinforcing feedback loop: blunted reward signals reduce exploratory transitions, yielding more predictable, low-information trajectories that further diminish intrinsic reward. The system can thus become self-maintaining even when the initial driver is not reward-related (e.g., cognitive load or inhibitory constraint), because reduced exploration progressively erodes intrinsic reinforcement.

Restoring the reward-exploration cycle requires interventions that enhance dopaminergic sensitivity and reintroduce cognitive novelty-seeking. Behavioral activation can increase contact with reinforcing experiences and environmental novelty (130), and cognitive-behavioral therapy (CBT) can introduce and train new cognitive strategies that promote exploratory engagement (131, 132). Pharmacological and neuromodulatory approaches that enhance reward sensitivity or reduce excessive constraint may likewise increase the hedonic value of cognitive progression, counteracting anhedonia. We further hypothesize that when thought becomes persistently

repetitive and low in informational gain, reduced intrinsic reward may contribute to anhedonia. Repetition itself is not inherently maladaptive; rather, FTP emphasizes trajectories that are low-gain and fail to update internal models. In this context, reduced discovery of novel associations may limit intrinsic reinforcement, sustain negative affect, and perpetuate rumination (116, 133, 134). This account generates testable predictions: individuals with greater thought rigidity should show reduced exploratory choice and blunted reward reactivity.

Plasticity and Cognitive Flexibility

At a cellular level, depression is characterized by reduced synaptic and dendritic plasticity across prefrontal and hippocampal regions (65, 66). Several accounts propose that these plasticity deficits may precede and amplify depressive symptoms, limiting the brain's capacity to update internal models or incorporate new emotional and cognitive information (65, 135). When neural circuits cannot readily modify synaptic weights or generate new dendritic connections, the formation of novel associative links is constrained, reducing the system's ability to sustain flexible cognitive trajectories. Reduced plasticity therefore contributes to affective inertia, cognitive rigidity, and potentially to the onset of depressive states themselves (136).

Pharmacological treatments that enhance plasticity provide causal support for this mechanism. Ketamine and serotonergic psychedelics induce rapid synaptogenesis and dendritic remodelling, opening short-lived "windows of plasticity" in which new cognitive and emotional associations can form (67). Similarly, selective serotonin reuptake inhibitors (SSRIs) have been shown to improve neuronal plasticity over longer timescales (65). Within the FTP framework, these interventions expand the brain's representational landscape, enabling more diverse and fluid thought trajectories.

These physiological observations converge with computational psychiatry models that describe depression as a low-variability regime, in which brain activity becomes overly predictable and transitions between neural states are infrequent (137). By enhancing plasticity, FTP-based interventions may shift the system back toward a flexible, metastable regime, restoring adaptive variability in both thought and mood.

The Facilitating Thought Progression (FTP) Framework: Mechanistic and Translational Implications

FTP as a Mechanistic Model for Mood Disorders

FTP functions as a translational model bridging basic neuroscience, computational psychiatry, and intervention science. It shifts emphasis from the *content* of thought to the *transitions* between mental representations, reframing depression as a disorder of state-space mobility. The core dimensions of cognitive progression map onto identifiable neural mechanisms: associative

breadth relates to large-scale network connectivity; thought speed reflects temporal properties of cortical dynamics and excitation-inhibition balance; cognitive flexibility depends on prefrontal-hippocampal connectivity dynamics and network reconfiguration; novelty processing engages dopaminergic mechanisms supporting exploratory cognition; and scope is governed by frontoparietal control over global vs. local integration.

When dysregulated, these processes collectively produce the cognitive rigidity and reward deficits characteristic of depression. Under this view, rumination is not merely a symptom but a mechanistic contributor that pushes cognition into a low-entropy, low-variability regime, restricting transitions across representational space.

Lastly, although FTP emphasizes thought dynamics over thought content, the two likely interact. Threat-related content may increase avoidance, narrow attention, and raise the perceived cost of exploration, thereby deepening attractor basins and reducing transition probabilities. In such cases, constrained dynamics may be partly driven by negative content that biases cognition toward safe, familiar trajectories.

Importantly, constrained dynamics can remain maladaptive even when content is not overtly negative if the system persistently fails to update models or integrate broader context. Even initially positive content may become problematic when cognition remains low in informational gain over time: novelty and learning signals decline, intrinsic reward diminishes, and repeated loops become more likely. Over time, this reward depletion may shift affective tone and motivation, linking seemingly benign content to depressive persistence through the same reward-exploration-rigidity feedback loop. FTP therefore treats content as a context-setting influence on trajectories, while dynamical parameters determine whether cognition can flexibly move out of those content-defined basins.

Translational Implications and Therapeutic Targets

Because FTP specifies the cognitive and neural mechanisms that support healthy thought flow, it highlights multiple intervention targets for restoring adaptive dynamics in depression. One therapeutic pathway involves reducing excessive inhibition and prefrontal hypercontrol. Treatments such as psilocybin, transcranial magnetic stimulation (TMS), or GABAergic modulation can transiently relax top-down constraints, increase neural entropy, and enable broader associative exploration (40, 96). A second avenue centers on enhancing reward responsiveness and novelty engagement. Pharmacological agents that boost dopaminergic or opioidergic signalling, as well as behavioral strategies that increase exposure to novelty, may re-engage the reward circuitry that reinforces exploratory thought (120, 121, 128, 129). Interventions that increase cortical plasticity offer another promising target. Glutamatergic agents such as ketamine and serotonergic psychedelics can rapidly enhance synaptogenesis and dendritic remodelling, potentially opening “windows of plasticity” during which new emotional

and cognitive associations become easier to form (65, 67). Such increases in plasticity expand the brain's representational landscape, supporting transitions out of rigid attractor states. Finally, exercises that increase associative breadth, semantic variability, or cognitive switching may restore more flexible trajectories through conceptual space. Recent clinical findings indeed show that cognitive interventions that directly train thought dynamics, and explicitly enhance thought progression can significantly reduce depressive symptoms (138). A more detailed comparison of FTP with existing models of depression and cognition is provided in the supplementary materials (see Supplementary S2). By decomposing cognitive rigidity or mental inertia into separable parameters of thought progression, FTP specifies which "lever" is altered and what behavioral and neural signature should follow (Fig. 3), enabling clearer cross-treatment comparison and mechanism-matched outcome selection.

Figure 3

Figure 3. FTP is a mechanistic bridge from neurocognitive levers to measurable thought dynamics: Top row: candidate mechanisms that constrain thought progression. Middle row: FTP dimensions of thought. Bottom row: observable behavioral signatures predicted by FTP. Lines illustrate the main hypothesized mappings between mechanisms, FTP dimensions, and measurement signatures.

Computational Psychiatry and Thought Dynamics

Recent advances in computational neuroscience (139, 140) and computational psychiatry (141-143) offer tools for formalizing the constructs central to FTP. Network control theory quantifies the brain's ability to transition between states and can reveal whether depression corresponds to greater control effort (144, 145). Complementing this systems-level view, models of spontaneous thought treat cognition as a dynamical process unfolding over an attractor landscape, enabling simulations of how excessive inhibition or reduced plasticity deepen attractor basins and stabilize ruminative loops [137]. Reinforcement-learning models further suggest that blunted reward sensitivity can bias cognition away from exploration, shifting thought trajectories toward exploitative, repetitive patterns consistent with reduced novelty-driven transitions (137, 141, 146).

In parallel, recent empirical work has begun to quantify thought dynamics directly. NLP-based semantic-trajectory analyses provide behavioral markers of cognitive flow such as semantic distance or variability derived from moment-to-moment thought reports (147, 148). These approaches can be combined with paradigms that sample ongoing experience (e.g., think-aloud sampling) and with dynamic neuroimaging analyses that identify latent brain states and their transition structure (26, 149, 150). Together, these methods enable cross-level mapping between observable properties of thought progression and neural state-transition signatures, providing a concrete empirical route for testing FTP's predictions (see Supplementary S3 for an extended 'testable predictions' section).

Conclusion

Depression can be understood as a disorder of impaired thought progression across five dimensions: breadth, speed, flexibility, novelty, and scope. FTP integrates evidence from cognitive science, computational psychiatry, and systems neuroscience to show how alterations in inhibition, reward processing, attractor stability, and neural plasticity bidirectionally shape cognitive flow and mood. In this view, rumination and negative mood reflect reduced cognitive fluidity, but the link is not one-way: promoting thought progression may itself help shift mood in an adaptive direction. FTP therefore offers a translational framework in which interventions that broaden, accelerate, or flexibilize thought, whether by reducing excessive inhibition, enhancing reward responsiveness, or increasing plasticity, may alleviate depressive symptoms. Future neuroimaging, computational, and longitudinal studies will be crucial for testing FTP and identifying its most informative markers.

Acknowledgments and Disclosures

M.B is supported by the Israeli Science Foundation (ISF grant #673/17) and the Sagol Family. S.B. is supported by the Edmond and Lily Safra Postdoctoral Fellowship and by a Minerva Fellowship of the Minerva Stiftung Gesellschaft fuer die Forschung mbH.

The authors report no biomedical financial interests or potential conflicts of interest.

Supplement Description:

Supplement Text and Table S1

Journal Pre-proof

References

1. Marx W, Penninx BW, Solmi M, Furukawa TA, Firth J, Carvalho AF, et al. Major depressive disorder. *Nature Reviews Disease Primers*. 2023;9(1):44.
2. Marvel CL, Paradiso S. Cognitive and neurological impairment in mood disorders. *Psychiatr Clin North Am*. 2004;27(1):19-36, vii-viii.
3. Lee AY, Sternthal B. The effects of positive mood on memory. *Journal of consumer research*. 1999;26(2):115-27.
4. Bower GH. Mood and memory. *American psychologist*. 1981;36(2):129.
5. Wadlinger HA, Isaacowitz DM. Positive mood broadens visual attention to positive stimuli. *Motivation and emotion*. 2006;30:87-99.
6. Taquet M, Quoidbach J, De Montjoye Y-A, Desseilles M, Gross JJ. Hedonism and the choice of everyday activities. *Proceedings of the national Academy of Sciences*. 2016;113(35):9769-73.
7. Forgas JP. Mood effects on decision making strategies. *Australian journal of Psychology*. 1989;41(2):197-214.
8. Shapiro D, Jamner LD, Goldstein IB, Delfino RJ. Striking a chord: Moods, blood pressure, and heart rate in everyday life. *Psychophysiology*. 2001;38(2):197-204.
9. Kok BE, Coffey KA, Cohn MA, Catalino LI, Vacharkulksemsuk T, Algoe SB, et al. How positive emotions build physical health: Perceived positive social connections account for the upward spiral between positive emotions and vagal tone. *Psychological science*. 2013;24(7):1123-32.
10. Patel R, Lloyd T, Jackson R, Ball M, Shetty H, Broadbent M, et al. Mood instability is a common feature of mental health disorders and is associated with poor clinical outcomes. *BMJ open*. 2015;5(5):e007504.
11. Abbott JT, Austerweil JL, Griffiths TL, editors. *Human memory search as a random walk in a semantic network*. NIPS; 2012: Citeseer.
12. Spinhoven P, van Hemert AM, Penninx BW. Repetitive negative thinking as a predictor of depression and anxiety: A longitudinal cohort study. *Journal of affective disorders*. 2018;241:216-25.
13. Nolen-Hoeksema S. The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *Journal of abnormal psychology*. 2000;109(3):504.
14. Beck AT. Thinking and depression: I. Idiosyncratic content and cognitive distortions. *Archives of general psychiatry*. 1963;9(4):324-33.
15. Northoff G, Hirjak D. Is depression a global brain disorder with topographic dynamic reorganization? *Translational Psychiatry*. 2024;14(1):278.
16. Braff DL, Beck AT. Thinking disorder in depression. *Archives of General Psychiatry*. 1974;31(4):456-9.
17. Ellis A. A sadly neglected cognitive element in depression. *Cognitive Therapy and Research*. 1987;11:121-45.
18. Teasdale JD, Segal ZV, Williams JMG, Ridgeway VA, Soulsby JM, Lau MA. Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *Journal of consulting and clinical psychology*. 2000;68(4):615.

19. Watkins E, Scott J, Wingrove J, Rimes K, Bathurst N, Steiner H, et al. Rumination-focused cognitive behaviour therapy for residual depression: A case series. *Behaviour research and therapy*. 2007;45(9):2144-54.
20. Kucyi A, Kam JWY, Andrews-Hanna JR, Christoff K, Whitfield-Gabrieli S. Recent advances in the neuroscience of spontaneous and off-task thought: implications for mental health. *Nat Ment Health*. 2023;1(11):827-40.
21. Migó M, Cooper JA, Kragel PA, Treadway MT. Spontaneous thought separates into clusters of negative, positive, and flexible thinking. *Communications Psychology*. 2025;3(1):21.
22. Fredrickson BL, Branigan C. Positive emotions broaden the scope of attention and thought-action repertoires. *Cognition & emotion*. 2005;19(3):313-32.
23. Isen AM, Daubman KA, Nowicki GP. Positive affect facilitates creative problem solving. *Journal of personality and social psychology*. 1987;52(6):1122.
24. Isen AM, Johnson M, Mertz E, Robinson GF. The influence of positive affect on the unusualness of word associations. *Journal of personality and social psychology*. 1985;48(6):1413.
25. Rowe G, Hirsh JB, Anderson AK. Positive affect increases the breadth of attentional selection. *Proceedings of the National Academy of Sciences*. 2007;104(1):383-8.
26. Raffaelli Q, Mills C, de Stefano N-A, Mehl MR, Chambers K, Fitzgerald SA, et al. The think aloud paradigm reveals differences in the content, dynamics and conceptual scope of resting state thought in trait brooding. *Scientific Reports*. 2021;11(1):19362.
27. Treynor W, Gonzalez R, Nolen-Hoeksema S. Rumination reconsidered: A psychometric analysis. *Cognitive therapy and research*. 2003;27:247-59.
28. Klinger E, Marchetti I, Koster EH. Spontaneous thought and goal pursuit: From functions such as planning to dysfunctions such as rumination. *The Oxford handbook of spontaneous thought: Mind-wandering, creativity, and dreaming*. 2018:215-47.
29. Robinson MS, Alloy LB. Negative cognitive styles and stress-reactive rumination interact to predict depression: A prospective study. *Cognitive Therapy and research*. 2003;27:275-91.
30. Lara ME, Klein DN, Kasch KL. Psychosocial predictors of the short-term course and outcome of major depression: a longitudinal study of a nonclinical sample with recent-onset episodes. *Journal of Abnormal Psychology*. 2000;109(4):644.
31. Dickson KS, Ciesla JA, Reilly LC. Rumination, worry, cognitive avoidance, and behavioral avoidance: Examination of temporal effects. *Behavior therapy*. 2012;43(3):629-40.
32. Akbari Chermahini S, Hommel B. Creative mood swings: divergent and convergent thinking affect mood in opposite ways. *Psychological research*. 2012;76:634-40.
33. Brunyé TT, Gagnon SA, Paczynski M, Shenhav A, Mahoney CR, Taylor HA. Happiness by association: Breadth of free association influences affective states. *Cognition*. 2013;127(1):93-8.
34. Mason MF, Bar M. The effect of mental progression on mood. *Journal of Experimental Psychology: General*. 2012;141(2):217.
35. LeMoult J, Yoon KL, Joormann J. Affective priming in major depressive disorder. *Frontiers in integrative neuroscience*. 2012;6:76.
36. Sass K, Habel U, Kellermann T, Mathiak K, Gauggel S, Kircher T. The influence of positive and negative emotional associations on semantic processing in depression: An fMRI study. *Human brain mapping*. 2014;35(2):471-82.

37. Besche-Richard C, Passerieux C, Hardy-Baylé M-C. Lexical decision tasks in depressive patients: semantic priming before and after clinical improvement. *European Psychiatry*. 2002;17(2):69-74.
38. Harmer CJ. Emotional processing and antidepressant action. *Behavioral neurobiology of depression and its treatment*. 2012:209-22.
39. Tranter R, Bell D, Gutting P, Harmer C, Healy D, Anderson IM. The effect of serotonergic and noradrenergic antidepressants on face emotion processing in depressed patients. *Journal of affective disorders*. 2009;118(1-3):87-93.
40. Daws RE, Timmermann C, Giribaldi B, Sexton JD, Wall MB, Erritzoe D, et al. Increased global integration in the brain after psilocybin therapy for depression. *Nature medicine*. 2022;28(4):844-51.
41. Northoff G. Beyond mood - depression as a speed disorder: biomarkers for abnormal slowness. *J Psychiatry Neurosci*. 2024;49(5):E357-E66.
42. Perini G, Cotta Ramusino M, Sinforiani E, Bernini S, Petrachi R, Costa A. Cognitive impairment in depression: recent advances and novel treatments. *Neuropsychiatric disease and treatment*. 2019:1249-58.
43. Rosser BA, Wright KA. The Impact of Thought Speed and Variability on Psychological State and Threat Perception: Further Exploration of the Theory of Mental Motion. *Cognitive Therapy and Research*. 2016;40:453-67.
44. Rostami S, Borjali A, Eskandari H, Rostami R, Scalabrini A, Northoff G. Slow and powerless thought dynamic relates to brooding in unipolar and bipolar depression. *Psychopathology*. 2022;55(5):258-72.
45. Den Hartog H, Derix M, Van Bemmelen A, Kremer B, Jolles J. Cognitive functioning in young and middle-aged unmedicated out-patients with major depression: testing the effort and cognitive speed hypotheses. *Psychological medicine*. 2003;33(8):1443-51.
46. Nuño L, Gómez-Benito J, Carmona VR, Pino O. A systematic review of executive function and information processing speed in major depression disorder. *Brain Sciences*. 2021;11(2):147.
47. Buyukdura JS, McClintock SM, Croarkin PE. Psychomotor retardation in depression: biological underpinnings, measurement, and treatment. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*. 2011;35(2):395-409.
48. Corfield EC, Martin NG, Nyholt DR. Co-occurrence and symptomatology of fatigue and depression. *Comprehensive psychiatry*. 2016;71:1-10.
49. Grahek I, Shenhav A, Musslick S, Krebs RM, Koster EH. Motivation and cognitive control in depression. *Neuroscience & Biobehavioral Reviews*. 2019;102:371-81.
50. Bar M. A cognitive neuroscience hypothesis of mood and depression. *Trends in cognitive sciences*. 2009;13(11):456-63.
51. Pronin E, Jacobs E, Wegner DM. Psychological effects of thought acceleration. *Emotion*. 2008;8(5):597.
52. Pronin E, Wegner DM. Manic thinking: Independent effects of thought speed and thought content on mood. *Psychological science*. 2006;17(9):807-13.
53. Yang K, Friedman-Wheeler DG, Pronin E. Thought acceleration boosts positive mood among individuals with minimal to moderate depressive symptoms. *Cognitive therapy and research*. 2014;38:261-9.

54. Channon S. Executive dysfunction in depression: the Wisconsin card sorting test. *Journal of affective disorders*. 1996;39(2):107-14.
55. Ashby FG, Isen AM. A neuropsychological theory of positive affect and its influence on cognition. *Psychological review*. 1999;106(3):529.
56. Baas M, De Dreu CK, Nijstad BA. A meta-analysis of 25 years of mood-creativity research: Hedonic tone, activation, or regulatory focus? *Psychological bulletin*. 2008;134(6):779.
57. Han W, Feng X, Zhang M, Peng K, Zhang D. Mood states and everyday creativity: Employing an experience sampling method and a day reconstruction method. *Frontiers in Psychology*. 2019;10:1698.
58. Stange JP, Alloy LB, Fresco DM. Inflexibility as a vulnerability to depression: A systematic qualitative review. *Clinical psychology: Science and practice*. 2017;24(3):245.
59. Stange JP, Connolly SL, Burke TA, Hamilton JL, Hamlat EJ, Abramson LY, et al. Inflexible cognition predicts first onset of major depressive episodes in adolescence. *Depression and anxiety*. 2016;33(11):1005-12.
60. Uddin LQ. Cognitive and behavioural flexibility: neural mechanisms and clinical considerations. *Nature Reviews Neuroscience*. 2021;22(3):167-79.
61. Dixon ML, De La Vega A, Mills C, Andrews-Hanna J, Spreng RN, Cole MW, et al. Heterogeneity within the frontoparietal control network and its relationship to the default and dorsal attention networks. *Proceedings of the National Academy of Sciences*. 2018;115(7):E1598-E607.
62. Zanto TP, Gazzaley A. Fronto-parietal network: flexible hub of cognitive control. *Trends in cognitive sciences*. 2013;17(12):602-3.
63. Spellman T, Rigotti M, Ahmari SE, Fusi S, Gogos JA, Gordon JA. Hippocampal–prefrontal input supports spatial encoding in working memory. *Nature*. 2015;522(7556):309-14.
64. Spellman T, Svei M, Kaminsky J, Manzano-Nieves G, Liston C. Prefrontal deep projection neurons enable cognitive flexibility via persistent feedback monitoring. *Cell*. 2021;184(10):2750-66. e17.
65. Duman RS, Aghajanian GK, Sanacora G, Krystal JH. Synaptic plasticity and depression: new insights from stress and rapid-acting antidepressants. *Nature medicine*. 2016;22(3):238-49.
66. Parekh PK, Johnson SB, Liston C. Synaptic mechanisms regulating mood state transitions in depression. *Annual Review of Neuroscience*. 2022;45(1):581-601.
67. Doss MK, Považan M, Rosenberg MD, Sepeda ND, Davis AK, Finan PH, et al. Psilocybin therapy increases cognitive and neural flexibility in patients with major depressive disorder. *Translational psychiatry*. 2021;11(1):574.
68. Lu JG, Akinola M, Mason MF. “Switching On” creativity: Task switching can increase creativity by reducing cognitive fixation. *Organizational Behavior and Human Decision Processes*. 2017;139:63-75.
69. Bruder GE, Kroppmann CJ, Kayser J, Stewart JW, McGrath PJ, Tenke CE. Reduced brain responses to novel sounds in depression: P3 findings in a novelty oddball task. *Psychiatry Research*. 2009;170(2-3):218-23.
70. Ivancovsky T, Baror S, Bar M. A shared novelty-seeking basis for creativity and curiosity: Response to the commentators. *Behavioral and Brain Sciences*. 2024;47:e119.

71. Smith R, Taylor S, Wilson RC, Chuning AE, Persich MR, Wang S, et al. Lower Levels of Directed Exploration and Reflective Thinking Are Associated With Greater Anxiety and Depression. *Front Psychiatry*. 2021;12:782136.
72. Bunzeck N, Doeller CF, Dolan RJ, Düzel E. Contextual interaction between novelty and reward processing within the mesolimbic system. *Human brain mapping*. 2012;33(6):1309-24.
73. Wittmann BC, Bunzeck N, Dolan RJ, Düzel E. Anticipation of novelty recruits reward system and hippocampus while promoting recollection. *Neuroimage*. 2007;38(1):194-202.
74. Craske MG, Dunn BD, Meuret AE, Rizvi SJ, Taylor CT. Positive affect and reward processing in the treatment of depression, anxiety and trauma. *Nature Reviews Psychology*. 2024;1-21.
75. Halahakoon DC, Kieslich K, O'Driscoll C, Nair A, Lewis G, Roiser JP. Reward-processing behavior in depressed participants relative to healthy volunteers: A systematic review and meta-analysis. *JAMA psychiatry*. 2020;77(12):1286-95.
76. Saragosa-Harris NM, Cohen AO, Reneau TR, Villano WJ, Heller AS, Hartley CA. Real-world exploration increases across adolescence and relates to affect, risk taking, and social connectivity. *Psychological Science*. 2022;33(10):1664-79.
77. Harel EV, Tennyson RL, Fava M, Bar M. Linking major depression and the neural substrates of associative processing. *Cognitive, Affective, & Behavioral Neuroscience*. 2016;16:1017-26.
78. Edwards JA, Weary G. Depression and the impression-formation continuum: piecemeal processing despite the availability of category information. *Journal of Personality and Social Psychology*. 1993;64(4):636.
79. Whitmer AJ, Gotlib IH. An attentional scope model of rumination. *Psychological bulletin*. 2013;139(5):1036.
80. Gasper K, Clore GL. Attending to the big picture: Mood and global versus local processing of visual information. *Psychological science*. 2002;13(1):34-40.
81. Ji L-J, Yap S, Best MW, McGeorge K. Global processing makes people happier than local processing. *Frontiers in Psychology*. 2019;10:670.
82. Jończyk R, Naranowicz M, Dębowska-Kozłowska K, Bromberek-Dyzman K. Negative mood constrains creative thinking in the native but not in the second language. *Thinking Skills and Creativity*. 2024;51:101457.
83. Bless H, Burger AM. Mood and the regulation of mental abstraction. *Current Directions in Psychological Science*. 2017;26(2):159-64.
84. Fredrickson BL. The broaden-and-build theory of positive emotions. *Philosophical transactions of the royal society of London Series B: Biological Sciences*. 2004;359(1449):1367-77.
85. Holmes AJ, Pizzagalli DA. Response conflict and frontocingulate dysfunction in unmedicated participants with major depression. *Neuropsychologia*. 2008;46(12):2904-13.
86. Keller AS, Leikauf JE, Holt-Gosselin B, Staveland BR, Williams LM. Paying attention to attention in depression. *Transl Psychiatry*. 2019;9(1):279.
87. Pizzagalli DA. Frontocingulate dysfunction in depression: toward biomarkers of treatment response. *Neuropsychopharmacology*. 2011;36(1):183-206.

88. Bennett MP, Knight R, Patel S, So T, Dunning D, Barnhofer T, et al. Decentering as a core component in the psychological treatment and prevention of youth anxiety and depression: a narrative review and insight report. *Translational Psychiatry*. 2021;11(1):288.
89. Travers-Hill E, Dunn BD, Hoppitt L, Hitchcock C, Dalgleish T. Beneficial effects of training in self-distancing and perspective broadening for people with a history of recurrent depression. *Behaviour Research and Therapy*. 2017;95:19-28.
90. Diener C, Kuehner C, Brusniak W, Ubl B, Wessa M, Flor H. A meta-analysis of neurofunctional imaging studies of emotion and cognition in major depression. *Neuroimage*. 2012;61(3):677-85.
91. Li Y, Liu X, Li J, Zhang Q, Pan N, Luo K, et al. Common and distinct patterns of brain activity alterations during inhibitory control in depression and psychostimulant users: a comparative meta-analysis of task-based fMRI studies. *Psychological medicine*. 2025;55:e218.
92. Zhou H-X, Chen X, Shen Y-Q, Li L, Chen N-X, Zhu Z-C, et al. Rumination and the default mode network: Meta-analysis of brain imaging studies and implications for depression. *Neuroimage*. 2020;206:116287.
93. Drevets WC, Videen TO, Price JL, Preskorn SH, Carmichael ST, Raichle ME. A functional anatomical study of unipolar depression. *Journal of Neuroscience*. 1992;12(9):3628-41.
94. Ressler KJ, Mayberg HS. Targeting abnormal neural circuits in mood and anxiety disorders: from the laboratory to the clinic. *Nature neuroscience*. 2007;10(9):1116-24.
95. LeDuke DO, Borio M, Miranda R, Tye KM. Anxiety and depression: A top-down, bottom-up model of circuit function. *Annals of the New York Academy of Sciences*. 2023;1525(1):70-87.
96. Carhart-Harris RL, Friston KJ. REBUS and the Anarchic Brain: Toward a Unified Model of the Brain Action of Psychedelics. *Pharmacol Rev*. 2019;71(3):316-44.
97. Barba T, Buehler S, Kettner H, Radu C, Cunha BG, Nutt DJ, et al. Effects of psilocybin versus escitalopram on rumination and thought suppression in depression. *BJPsych Open*. 2022;8(5):e163.
98. Beaty RE, Kenett YN. Associative thinking at the core of creativity. *Trends in cognitive sciences*. 2023;27(7):671-83.
99. Koivisto M, Toivanen H. Schizotypy and Creativity: Divergent Thinking, Inhibitory Control, and the Spontaneous Flow of Thought. *Creativity Research Journal*. 2024:1-21.
100. Moritz S, Mersmann K, Kloss M, Jacobsen D, Wilke U, Andresen B, et al. 'Hyper-priming' in thought-disordered schizophrenic patients. *Psychological medicine*. 2001;31(2):221-9.
101. Miller R. Hyperactivity of associations in psychosis. *Australian & New Zealand Journal of Psychiatry*. 1989;23(2):241-8.
102. De Lissnyder E, Koster EH, Derakshan N, De Raedt R. The association between depressive symptoms and executive control impairments in response to emotional and non-emotional information. *Cognition and Emotion*. 2010;24(2):264-80.
103. Yoon KL, LeMoult J, Joormann J. Updating emotional content in working memory: a depression-specific deficit? *Journal of Behavior Therapy and Experimental Psychiatry*. 2014;45(3):368-74.
104. Gotlib IH, Joormann J. Cognition and depression: current status and future directions. *Annu Rev Clin Psychol*. 2010;6:285-312.

105. Vonasch AJ, Vohs KD, Pocheptsova Ghosh A, Baumeister RF. Ego depletion induces mental passivity: Behavioral effects beyond impulse control. *Motivation Science*. 2017;3(4):321.
106. Baror S, Bar M. Increased associative interference under high cognitive load. *Scientific reports*. 2022;12(1):1766.
107. Baror S, Bar M. Associative activation and its relation to exploration and exploitation in the brain. *Psychological science*. 2016;27(6):776-89.
108. Brownlow JA, Miller KE, Gehrman PR. Insomnia and cognitive performance. *Sleep medicine clinics*. 2019;15(1):71.
109. Olivera-López C, Jiménez-Genchi A, Ortega-Robles D, Valencia-Flores M, Cansino S, Salvador-Cruz J. Polysomnographic parameters associated with cognitive function in patients with major depression and insomnia. *CNS spectrums*. 2024;29(3):197-205.
110. Lim J, Dinges DF. A meta-analysis of the impact of short-term sleep deprivation on cognitive variables. *Psychological bulletin*. 2010;136(3):375.
111. Massar SA, Lim J, Sasmita K, Chee MW. Sleep deprivation increases the costs of attentional effort: Performance, preference and pupil size. *Neuropsychologia*. 2019;123:169-77.
112. Westgate EC, Wilson TD, Buttrick NR, Furrer RA, Gilbert DT. What makes thinking for pleasure pleasurable? *Emotion*. 2021;21(5):981.
113. Barrett LF. The theory of constructed emotion: an active inference account of interoception and categorization. *Social cognitive and affective neuroscience*. 2017;12(1):1-23.
114. Kube T, Kirchner L, Gärtner T, Glombiewski JA. How negative mood hinders belief updating in depression: results from two experimental studies. *Psychological medicine*. 2023;53(4):1288-301.
115. Herz N, Baror S, Bar M. Overarching states of mind. *Trends in Cognitive Sciences*. 2020;24(3):184-99.
116. Admon R, Pizzagalli DA. Dysfunctional reward processing in depression. *Current opinion in psychology*. 2015;4:114-8.
117. Biederman I, Vessel EA. Perceptual pleasure and the brain: A novel theory explains why the brain craves information and seeks it through the senses. *American scientist*. 2006;94(3):247-53.
118. Kenett YN, Humphries S, Chatterjee A. A thirst for knowledge: Grounding curiosity, creativity, and aesthetics in memory and reward neural systems. *Creativity Research Journal*. 2023;35(3):412-26.
119. Koeppe MJ, Hammers A, Lawrence AD, Asselin M-C, Grasby PM, Bench C. Evidence for endogenous opioid release in the amygdala during positive emotion. *Neuroimage*. 2009;44(1):252-6.
120. Gruber MJ, Gelman BD, Ranganath C. States of curiosity modulate hippocampus-dependent learning via the dopaminergic circuit. *Neuron*. 2014;84(2):486-96.
121. Kang MJ, Hsu M, Krajbich IM, Loewenstein G, McClure SM, Wang JT-y, et al. The wick in the candle of learning: Epistemic curiosity activates reward circuitry and enhances memory. *Psychological science*. 2009;20(8):963-73.
122. Trapp S, Shenhav A, Bitzer S, Bar M. Human preferences are biased towards associative information. *Cognition and Emotion*. 2015;29(6):1054-68.
123. Bar M. *Mindwandering: How It Can Improve Your Mood and Boost Your Creativity*: Bloomsbury Publishing; 2022.

124. Briellmann AA, Pelli DG. Beauty requires thought. *Current Biology*. 2017;27(10):1506-13. e3.
125. O'Brien E. Enjoy it again: Repeat experiences are less repetitive than people think. *Journal of Personality and Social Psychology*. 2019;116(4):519.
126. Zuckerman M. Sensation seeking (psychology revivals): Beyond the optimal level of arousal: Psychology Press; 2014.
127. Lewis ME, Mishkin M, Bragin E, Brown RM, Pert CB, Pert A. Opiate receptor gradients in monkey cerebral cortex: Correspondence with sensory processing hierarchies. *Science*. 1981;211(4487):1166-9.
128. Jelen LA, Stone JM, Young AH, Mehta MA. The opioid system in depression. *Neuroscience & Biobehavioral Reviews*. 2022;140:104800.
129. Addicott MA, Pearson JM, Sweitzer MM, Barack DL, Platt ML. A primer on foraging and the explore/exploit trade-off for psychiatry research. *Neuropsychopharmacology*. 2017;42(10):1931-9.
130. Cuijpers P, Karyotaki E, Harrer M, Stikkelbroek Y. Individual behavioral activation in the treatment of depression: A meta analysis. *Psychother Res*. 2023;33(7):886-97.
131. Norbury A, Hauser TU, Fleming SM, Dolan RJ, Huys QJ. Different components of cognitive-behavioral therapy affect specific cognitive mechanisms. *Science Advances*. 2024;10(13):eadk3222.
132. Yang Z, Oathes DJ, Linn KA, Bruce SE, Satterthwaite TD, Cook PA, et al. Cognitive behavioral therapy is associated with enhanced cognitive control network activity in major depression and posttraumatic stress disorder. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. 2018;3(4):311-9.
133. Nusslock R, Alloy LB. Reward processing and mood-related symptoms: An RDoC and translational neuroscience perspective. *Journal of affective disorders*. 2017;216:3-16.
134. Rutherford AV, McDougle SD, Joormann J. "Don't [ruminate], be happy": A cognitive perspective linking depression and anhedonia. *Clinical Psychology Review*. 2023;101:102255.
135. Liu W, Ge T, Leng Y, Pan Z, Fan J, Yang W, et al. The role of neural plasticity in depression: from hippocampus to prefrontal cortex. *Neural plasticity*. 2017;2017(1):6871089.
136. Maramis MM, Mahajudin MS, Khotib J. Impaired cognitive flexibility and working memory precedes depression: a rat model to study depression. *Neuropsychobiology*. 2021;80(3):225-33.
137. Rolls ET. Attractor cortical neurodynamics, schizophrenia, and depression. *Transl Psychiatry*. 2021;11(1):215.
138. Yatziv S-L, Pedrelli P, Baror S, DeCaro SA, Shachar N, Sofer B, et al. Facilitating Thought Progression to Reduce Depressive Symptoms: Randomized Controlled Trial. *Journal of Medical Internet Research*. 2024;26:e56201.
139. Kriegeskorte N, Douglas PK. Cognitive computational neuroscience. *Nat Neurosci*. 2018;21(9):1148-60.
140. Kringelbach ML, Deco G. Brain States and Transitions: Insights from Computational Neuroscience. *Cell Rep*. 2020;32(10):108128.
141. Huys QJ, Maia TV, Frank MJ. Computational psychiatry as a bridge from neuroscience to clinical applications. *Nat Neurosci*. 2016;19(3):404-13.

142. Khaleghi A, Mohammadi MR, Shahi K, Nasrabadi AM. Computational Neuroscience Approach to Psychiatry: A Review on Theory-driven Approaches. *Clin Psychopharmacol Neurosci*. 2022;20(1):26-36.
143. Wang XJ, Krystal JH. Computational psychiatry. *Neuron*. 2014;84(3):638-54.
144. Clark JE, Watson S. Modelling mood updating: a proof of principle study. *Br J Psychiatry*. 2023;222(3):125-34.
145. Kenett YN, Beaty RE, Medaglia JD. A computational network control theory analysis of depression symptoms. *Personal Neurosci*. 2018;1.
146. Chen C, Takahashi T, Nakagawa S, Inoue T, Kusumi I. Reinforcement learning in depression: A review of computational research. *Neurosci Biobehav Rev*. 2015;55:247-67.
147. Li HX, Chen X, Wang ZH, Lu B, Liao YF, Li XY, et al. Characterizing human spontaneous thoughts and its application in major depressive disorder. *J Affect Disord*. 2024;365:276-84.
148. Palaniyappan L, Wang YL, Meister F. Disorganisation and depression: a re-examination of how we think and speak when depressed. *Eur Arch Psychiatry Clin Neurosci*. 2025.
149. Mildner JN, Tamir DI. Why do we think? The dynamics of spontaneous thought reveal its functions. *PNAS nexus*. 2024;3(6):pgae230.
150. Yu Y, Raffaelli Q, Freveletti D, Brossart J, Andrews E, Wilcox RR, et al. Mapping Content and Dynamics in the Stream of Consciousness: a Latent Brain State Analysis.

